

Reveticiclina F®
Reg. Key SAGARPA Q-0524-039*
FOR VETERINARY USE

Penicillin-Streptomycin-Dexamethasone

Injectable broad-spectrum antibiotic powder and diluent reconstitution

DRUG DESCRIPTION

Every mL of reconstituted product contains

Penicillin G Procaine 150,000 IU

Penicillin G Sodium 50,000 IU

Streptomycin sulfate, equivalent to:

Streptomycin base 0.25 g

Dexamethasone 0.25 mg

Dipyrone 200 mg

Guaiacol 10 mg

Excipient c.b.p. 1 mL

Reveticiclina® F is a reconstitution injectable broad-spectrum bactericidal antibiotic antipruritic and expectorant effective against Gram positive and Gram negative bacteria, the penicillins contained in this product belong to the Beta-lactamases group, the Streptomycin is included as an aminoglycoside and the Dexamethasone included belongs to the glucocorticoids group.

Reveticiclina® F is recommended as treatment of infections such as: mastitis, metritis, pneumonia, rhinitis and bacterial arthritis in Dairy Cattle, sheep, goats, swine, horses, cats and dogs. For barnyard fowl is recommended as treatment of ulcerative enteritis, borreliosis, psittacosis, colibacillosis and avian cholera.

Reveticiclina® F has a synergistic effect among its antibiotics which shows more effectiveness; it eliminates bacterial strains that are resistant to other medication.

Reveticiclina® F Due to its antiphlogistic effect it works directly on the capillary vasodilation in swollen conditions, accelerating the tissues' process of involution and allowing a better penetration of the antibiotics into the spot of the infection.

DRUG USE

Broad-spectrum bactericidal antibiotic with anti-inflammatory effect indicated for treatment of infections caused by bacteria sensitive to the formula.

DOSAGE Cats: 0.25 to 0.5 mL (total dose)

Dogs: 0.5 to 3 mL (total dose)

Swine, sheep and goats: 1 to 2 mL per 20 kg.

Dairy Cattle y horses: 1 mL per 20 kg.

Frequency: In any case every 12 to 24 hours according to prescription.

ROUTE OF ADMINISTRATION

Intramuscular.

DRUG WARNINGS

This product should not be used during gestation period due to the effect of the dexamethasone on the placental. Do not use on animals with hypersensitivity to any of the components of the formula. Stop use immediately in case of hypersensitivity reactions to medication and treat with antihistamines. Empty containers and any residue must be disposed of by burying or incinerating. For reconstitution use the whole content in the vial and the included diluent. Use the product immediately after reconstitution.

Keep this product in a cool and dry place protected from light at a temperature between 15° and 30 °C Do not use in horses destined for human consumption.

Keep out of reach of children and pets.

WITHDRAWAL PERIOD

Milk withdrawal: 3 days

Meat withdrawal: 30 days

Consult a veterinarian

Prescription Only Medicine – Veterinarian (POM-V category)

PRESENTATION

1 vial x 10 mL (2 000 000 I.U.)

1 vial x 20 mL (4 000 000 I.U.)

1 vial x 30 mL (6 000 000 I.U.)

REGISTRATION KEYS

Costa Rica MAG MX7-05-05-151

El Salvador VET.96-7-1190

Honduras PF-915

Nicaragua 2378

Dominican Rep. 2191-95

MADE IN MEXICO BY: RIVERFARMA, S.A. DE C.V. □ Prol. Calz. de La Viga No. 1949, Col. Prado Churubusco, Del. Coyoacán. C.P.04230, México, D.F. E-mail: riverfarma@riverfarma.com.mx Web: www.riverfarma.com.mx

Tel. 6389-6687, 6389-6688, 6389-6689. Within Mexico Toll Free: 01-800-505-7777