

Analvet®
Reg. Key SAGARPA Q-0524-017*
FOR VETERINARY USE
Sodium Dipyron
Injectable Solution

Antipruritic, anti-inflammatory, antispasmodic, non steroidal analgesic

DRUG DESCRIPTION

Each 100 mL contain Sodium Dipyron.....50 g

Excipient c.b.p.100 ml

Analvet® is an analgesic that works at the thalamus level at the cerebral cortex inhibiting the synthesis of prostaglandins (histamine and bradykinin) responsible for any painful affection. As main components it contains dipyron and metamizole which belong to the pyrazolone group with pain killing effects. Non- narcotic, antipyretic, anti-inflammatory and antispasmodic.

- It works on the thermoregulatory central and peripheral systems in order to rapidly reduce fever of infectious diseases such as: pneumonic pasteurellosis, other pneumonias, piroplasmosis, anaplasmosis and salmonellosis.
- In horses it helps as treatment of spasms and excessive intestinal peristalsis during colics. In spasmodic conditions of the uterine musculature during birth of mammals. In swine it is particularly useful to avoid that first time moms devour their piglets (30 mL intramuscular route).
- It works as anti-inflammatory aid (traumatism, surgical wound care, arthritis, tendonitis, myositis, rheumatism and lumbago in horses).

DRUG USE

Recommended for symptomatic traumatism, neuralgic affections and postoperative pain. This medication also has an antipyretic effect that helps reduce fever and colics in horses.

DOSAGE

Horses: 4 to 8 mL per 100 kg

Dairy Cattle, sheep, goats and swine: 1 mL per 10 kg.

Dogs and Cats: .1 mL per 2 kg In any case administer every 8 hours for no more than 5 days.

ROUTE OF ADMINISTRATION

Slow intravenous or deep intramuscular. **DRUG WARNINGS** In case of over dose or long-lasting treatment it can cause moderate hepatic affection. Do not mix with chlorpromazine, barbiturates or phenylbutazone as it may cause severe hypothermia. Not recommended for animals with blood dyscrasia. Do not use in horses destined for human consumption.

WITHDRAWAL PERIOD

Meat withdrawal: 24 hours

Milk withdrawal: 24 hours

Empty containers and any residue must be disposed of by burying or incinerating.

Keep this product in a cool and dry place protected from light at a temperature between 15° and 30 °C

Keep out of reach of children and pets.

Consult a veterinarian

Prescription Only Medicine – Veterinarian (POM-V category) PRESENTATION 50 or 100 mL vial

REGISTRATION KEYS

Costa Rica MAG MX7-02-03-056

El Salvador VE2013094811

Honduras PF-836

Nicaragua 3814

Panama 1491-92

DominicanRep. RF-557-11

MADE IN MEXICO BY: RIVERFARMA, S.A. DE C.V.

Prol. Calz. de La Viga No. 1949, Col. Prado Churubusco, Del. Coyoacán. C.P.04230, México, D.F.

E-mail: riverfarma@riverfarma.com.mx Web: www.riverfarma.com.mx Tel. 6389-6687, 6389-6688, 6389-6689. Within Mexico Toll Free: 01-800-505-7777