

Micofin® 1%
Reg. Key SAGARPA Q-0524-103*
FOR VETERINARY USE
Terbinafine
Antimycotic topical solution

DRUG DESCRIPTION

Every 100 mL contain
Terbinafine Clorhydrate Equivalent to 1 g of terbinafine base
Excipient c.b.p. 100 mL

Micofin 1% is a broad-spectrum antimycotic topical solution. Its active component, terbinafine chemically belongs to the allylamines

Micofin 1% is recommended for treating skin micotic infections (ringworm) and claws caused by dermatophytes (*Microsporum* spp., *Trichophyton* spp and *Epidermophyton floccosum*) as well as infections caused by yeast such as *Candida* (*Candida albicans*), in dogs and cats.

Micofin 1% has a fungicidal effect even in low concentrations.

Micofin 1% achieves high concentrations on the affected spots even though they might be difficult to reach like skin, claws and hair due to its keratin.

Micofin 1% Due to its effectiveness it's first choice antimycotic treatment for superficial mycoses, it acts immediately and its effective from the beginning.

Micofin 1% it is more effective than Imidazoles and griseofulvina which implies less number of relapses and treatment period reductions.

Micofin 1% Keeps under control itchiness and nibbling on the affected areas, stops hair loss due to fungus, it also improves pets' physical shape and aspect.

Micofin 1% It can be used along antibiotics (Enrocilina® Tablets) and acaricides (Endovet® Tablets, Endovet®-Ces Tablets and/or Endovet®-Ces Injectable) when treating dermatitis.

DRUG USE

It is recommended for treating skin micotic infections (ringworm) and claws caused by dermatophytes (*Microsporum* spp., *Trichophyton* spp and *Epidermophyton floccosum*) as well as infections caused by yeast such as *Candida* (*Candida albicans*), in dogs and cats.

DOSAGE

Apply a light layer on affected skin and nearby areas, massage gently with gauze and wear latex gloves. Frequency: Apply every 12 to 24 hours during 1 to 2 weeks depending on the severity of the case and the extension of the infection. Treating mycosis the time required ranges from 6 weeks to 3 months.

ROUTE OF ADMINISTRATION

Topical

DRUG WARNINGS

Wear medical gloves for applying

For external use only Avoid contact with eyes Clean and dry affected spots before applying Irregular or interrupted use lead to relapse, if there are no improving signs after 2 weeks with the treatment check the diagnosis The total healing could be achieved after many weeks from mycotic healing. Empty containers and any residue must be disposed of by burying or incinerating.

Keep this product in a cool and dry place protected from light at a temperature between 15° and 30°C

Keep out of reach of children and pets.

Consult a veterinarian Prescription Only Medicine – Veterinarian (POM-V category)

PRESENTATION

30 mL vial

REGISTRATION KEYS

Costa Rica MAG MX-17-10-3168

El Salvador VE2013094813

Honduras PF-3588

Nicaragua 6741

Panama RF-3343-05

Dominican Rep. 6557

MADE IN MEXICO BY: RIVERFARMA, S.A. DE C.V. Prol. Calz. de La Viga No. 1949, Col. Prado Churubusco, Del. Coyoacán. C.P.04230, México, D.F. E-mail: riverfarma@riverfarma.com.mx Web: www.riverfarma.com.mx

Tel. 6389-6687, 6389-6688, 6389-6689. Within Mexico Toll Free: 01-800-505-7777