

Sulfatrim
Reg. Key SAGARPA Q-0524-013*
FOR VETERINARY USE
Trimethoprim - Sulfamethoxazole
Injectable Solution

Broad-spectrum antimicrobial bactericidal injectable solution

DRUG DESCRIPTION

Every 100 mL contain

Trimethoprim	4 g
Sulfamethoxazole	20 g
Excipient c.b.p.	100 ml

Sulfatrim® is a broad-spectrum antimicrobial bactericidal antibiotic injectable solution that contains sulfamethoxazole (part of the sulfonamide group) and trimethoprim (a diaminopyridine).

Sulfatrim® is recommended for treatment of infections of the gastrointestinal tract, urogenital system, respiratory tract and septicemias caused by Gram positive, Gram negative bacteria and protozoa. The use is recommended in Dairy Cattle, sheep, goats, swine, dogs, cats and barnyard fowl.

Sulfatrim® Thanks to its double inhibitor effect on bacteria it works simultaneously on the biosynthesis of folic acid showing a broad-spectrum and decreasing the risk of resistance development, acting even against those microorganisms that are resistant to other medications.

Sulfatrim® achieves high tissue and blood concentrations that produce an immediate clinic response and higher effectiveness.

Sulfatrim® has a broader spectrum compared to other medications such as Beta- lactamases, aminoglycosides and some macrolides.

Sulfatrim® is an effective treatment against coccidiosis.

DRUG USE

For treatment of infections of the gastrointestinal tract, urogenital system, respiratory tract and septicemias.

DOSAGE

The period and frequency of consumption must be adjusted according to the animal. The average recommended dose is 15 to 20 mg/kg or 1 mL per 12 to 16 kg.

Dairy Cattle and adult horses	20 to 30 mL
Calves and colts	5 to 15 mL

Sheep and goats	3 to 5 mL
Lambs and suckling pigs	0.5 to 1 mL
Adult swine	5 to 10 mL
Dogs and Cats	1 to 3 mL
Lightweight Birds	0.2 mL per kg
Heavyweight Birds	0.15 mL per kg

Oral administration: 1 mL of Sulfatrim per .5 L of water.

Depending on the severity of the infection it is recommended to repeat the dose every 24 to 48 hours.

ROUTE OF ADMINISTRATION

Intravenous, intramuscular, subcutaneous or oral.

DRUG WARNINGS

A sulfanomide swift intravenous injection may cause respiratory problems and collapse, that's why slow injection is advised. The incorrect intravenous administration technique of Sulfatrim may cause serious perivascular reactions and thrombophlebitis. Do not mix Sulfatrim with other products in the same syringe. Do not administer to animals with serious hepatic affections, blood dyscrasia, renal insufficiency or hypersensitivity to the formula. In case of overdose proper ventilation is recommended as well as treating with intravenous electrolyte solution.

Do not use in horses destined for human consumption.

WITHDRAWAL PERIOD

Milk withdrawal: 5 days

Poultry withdrawal: 5 days

Egg withdrawal: 8 days

Meat withdrawal: 12 days

Keep this product in a cool and dry place protected from light at a temperature between 15° and 30 °C

Keep out of reach of children and pets. Consult a veterinarian Prescription Only Medicine – Veterinarian (POM-V category)

PRESENTATION

10, 25, 50 and 100 mL vials

REGISTRATION KEYS

Costa Rica MAG MX7-46-08-944

El Salvador VE2013094789

Nicaragua 2379

MADE IN MEXICO BY: RIVERFARMA, S.A. DE C.V. Prol. Calz. de La Viga No. 1949, Col. Prado Churubusco, Del. Coyoacán. C.P.04230, México, D.F. E-mail: riverfarma@riverfarma.com.mx Web: www.riverfarma.com.mx

Tel. 6389-6687, 6389-6688, 6389-6689. Within Mexico Toll Free: 01-800-505-7777